

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.

THIS PAGE BLANK (USPTO)



The
Patent
Office

PCT

90 / 00526	
REC'D	18 MAR 1998
WIPO	PCT

4

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP9 1RH

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

PRIORITY DOCUMENT

Signed

Dated 2 February 1998



THIS PAGE BLANK (USPTO)

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

27JAN97 E248916-1 C03182
P01/7700 25.00 The Patent Office

Cardiff Road
Newport
Gwent NP9 1RH

25 JAN 1997

1. Your reference

CV-0244 GB

25 JAN 1997

2. Patent application number

(The Patent Office will fill in this part)

9701552.3

3. Full name, address and postcode of the or of each applicant (underline all surnames)

Bristol-Myers Squibb Company
345 Park Avenue
New York
New York 10154
United States of America
Incorporated in Delaware

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation

44-45882001

4. Title of the invention

MULTI-DOSE WOUND GEL

5. Name of your agent (if you have one)

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

Julie MAYS
Bristol-Myers Squibb Company
Patent Department
Swakeleys House
Milton Road
Ickenham, Uxbridge UB10 8NS

Patents ADP number (if you know it)

64591163002

642246002

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number
(if you know it)

Date of filing
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
 - b) there is an inventor who is not named as an applicant, or
 - c) any named applicant is a corporate body.
- See note (d))

Yes

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description 10
Claim(s) 1
Abstract 0
Drawing(s) 1

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77) 1

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

11.

I/We request the grant of a patent on the basis of this application.

Signature

Date 24 Jan 1977

12. Name and daytime telephone number of person to contact in the United Kingdom

Julie MAYS

01895 628209

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

5

MULTI-DOSE WOUND GEL

This invention relates to a multi-dose wound gel. More particularly, this invention relates to a wound gel packaged in a multi-dose container, useful for treating
10 wounds.

It is well known that the cleansing and debriding of wounds and the removal of wound exudate is important to the process of healing wounds. Commonly used wound dressings comprise gauze, foams, sponges, cotton wads or other
15 fibrous materials. Gauze and other fibrous materials absorb fluids by capillary action with the disadvantage that when new tissue is formed as part of the healing process, it engulfs the fibres and is torn when the material is removed causing wound injury.

20 Various other materials have been used, such as gels hydrogels, granules and pastes to remove exudates from wounds. Certain wound gels are known to promote the healing of wounds. For instance they can keep the wound bed moist, cleanse the wound, debride necrotic matter by
25 fluid donation and absorb exudate. Freshly generated tissue does not grow into the gel and thus injury on removal is avoided.

The gels are usually packaged in a tube and applied to the wound from the tube. The gels are usually in either a
30 sterile or a preserved state. If packaged in a multi-dose tube there is a risk with some gels that once the tube is

5 opened bacteria will enter the tube and proliferate in the
gel. For this reason some manufacturers include
preservatives in the gel or package in single dose tubes.
Some health care professionals are reluctant to introduce
preservatives to a wound and so use single dose tubes
10 containing sterile gel. This adds to the cost of the
product and results in wastage if the whole contents of the
tube is not used. There is therefore a need for a multi-
dose gel packaged in such a way that contamination is
minimised once the packaging is opened.

15 We have now found that is possible to package a gel in
multi-dose packaging which minimises contamination once
opened. This is achieved by the use of a barrier aerosol.

Accordingly the invention provides a barrier aerosol vessel
containing a wound gel for the treatment of wounds.

20 Aerosol barrier vessels are of the type where the product
to be dispensed and the pressure generating media, ie the
propellant, are maintained in isolation through separation
on opposite sides of a barrier. This has many advantages
in the context of wound gels. Firstly, because there is
25 positive pressure in the container, the vessel can be made
to be self-sealing. This aids maintenance of product
sterility. Secondly, it is possible to use an aerosol
single-handed which makes application of the gel to the
wound particularly easy. In the case of sinus wounds,
30 where there is undermining of the tissue beneath the wound
site and beyond its surface periphery, it is possible to
insert the nozzle through the sinus to fill the cavity.

5 Three main variants of barrier vessel exist. In a piston-type barrier vessel the barrier is a piston-like component that is mounted in the container in sliding relation to the inside surface of the container. The product to be dispensed is disposed on the valved side of the piston and
10 the propellant, which generates pressure within the container, is on the opposite side of the piston. Examples of piston-type barrier packs are described in US 3,033,923, 3,756,476 and 3,929,132.

In a second variant of an aerosol barrier vessel, a
15 flexible collapsible inner container is affixed within an outer container opening either to the aerosol discharge valve or to the bead of the container opening. Patents which illustrate a barrier vessel of this variant are described in US 3,788,521, 3,896,970 and 4,067,499.

20 In a third variant the barrier vessel is an unfolding cup-shaped barrier wherein the barrier has an outer wall terminating in a sealing flange, said outer wall being disposed contiguous to the inner wall of the container. The inner wall of the barrier is initially folded within
25 the outer wall, the inner wall terminating in an end closing portion. The barrier is contained in a valved aerosol container and sealed at the joint formed between the sidewall and the bottom end closure of the container. Product is admitted through the valved opening of the
30 container and propellant through a port in the bottom end closure of the container. Actuation of the valve reduces the pressure in the product compartment and results in the inner wall of the barrier unfolding from within the outer

5 wall of the barrier and causing the end-closing portion of the inner wall of the barrier to advance and thereby urge the product toward the discharge port. This type of barrier vessel is illustrated in US 3,109,463 and WO 96/02439.

10 The barrier aerosol vessel preferably used in the present invention is of the second or third type and comprises an inner container which contains the gel sealed by an opening valve with a discharge port for discharging the gel, an
15 outer casing container covering the inner container and a pressure medium interposed between the inner container and the outer casing container. The use of this type of container enables the inner container to be filled with non-sterile gel while assembled in the outer casing container, sealed by the valve and then sterilised by steam
20 sterilisation. The pressure medium can then be introduced without compromising the sterility of the product.

Preferably the inner container is made of a thin flexible material such as plastic or metal foil, although metal foil is especially preferred to maintain sterility if a sterile
25 gel is used. The outer casing container is also preferably metal such as aluminium which is pressure resistant. The outer container is preferably formed by compression moulding, thermoforming or the like, the inside provided with an inner protective coating and primed and the base
30 provided with a valve to enable the container to be pressurised once the inner container has been filled and sealed. The inner container is preferably sealed by a valve which comprises a cup and a discharge port and closes

5 off the outer container.

A barrier vessel suitable for use in the present invention is illustrated in the following figures:

Figure 1 is an elevation in section of one embodiment of the invention.

10 Figure 2 is a perspective view of the vessel of the invention in use.

An example of a barrier aerosol vessel (2) suitable for use in the present invention is shown in Figure 1 and comprises an inner container (4) which contains a gel (6) sealed by
15 an opening valve (8) for discharging the gel (6), an outer casing container (10) covering the inner container (4) and a pressure medium (12) interposed between the inner container (4) and the outer casing container (10). The
20 outer casing container is provided with a sealable port (14) to enable the pressure medium (12) to be introduced. The opening valve (8) comprises a cup (16) and discharge port (18). The whole of the opening valve (8), including cup (16) and port (18) will conventionally be covered with an applicator (not shown). Depression of which by the user
25 causes the gel to exit the port (18) into a conventional nozzle or an extension nozzle depending on the use. Such nozzles can be separately packaged for single use.

Figure 2 shows the aerosol vessel of the invention in use. In this view an applicator has been placed on the opening
30 valve to aid application of the gel to a wound.

5 When the gel is to be dispensed the valve (8) is actuated, the pressure medium acts to collapse the inner container (4) and gel (6) flows from the discharge port (18) of valve (8).

The gel for use in the present invention is preferably a
10 hydrocolloid gel and comprises a cellulose derivative, water and a polyol component. Such gels are described in EP-A-576523. More preferred is a gel comprising:

- (a) from about 0.05% to 10% by weight of a natural gelling agent;
- 15 (b) from about 1.0% to 10% by weight of a hydrocolloid;
- (c) from about 5.0% to 30.0% by weight of a glycol and
- (d) at least 50% by weight of water.

The most preferred composition contains 0.1% pectin, 3.4% sodium carboxymethyl cellulose, 15% propylene glycol and
20 81.5% water. Such gels are described in EP-A-567311 and EP-A-666081.

The natural gelling agent is preferably selected from pectin, alginic acid and salts thereof, carageenan, tragacanth, acacia, locust bean gum, guar gum, starch, agar
25 and gelatin. More preferably the pectin is pectin with a high ester content derived from citrus peel consisting chiefly of the partial methyl esters of polygalachronic acid (approximately 65% of the carboxyl groups are esterified). Representatives of the pectin useful in the
30 gel composition is that marketed under the name GENU pectin type VIS-L by Copenhagen Pectin. The natural gelling agent is preferably present in an amount from 0.05% to 1.0% by

5 weight.

The hydrocolloid is preferably selected from sodium carboxymethyl cellulose, methyl cellulose, hydroxypropylmethyl cellulose, hydroxyethyl cellulose, hydroxymethylcellulose, hydroxypropyl cellulose, 10 carboxyvinyl polymer and salts thereof, poloxamer for example Pluronic F127, xanthan gum, povidone, modified starches, and guar derivatives. The carboxymethyl cellulose is preferably sodium carboxymethyl cellulose present in an amount from about 2.0% to 4.5% by weight. The preferred 15 sodium carboxymethyl cellulose is a high viscosity sodium carboxymethyl cellulose (typically in the range 2000 - 4500 cps as measured by Brookfield LV Viscometry of a 1% solution, oven dry basis, 25°C and spindle 4/30 rpm.

The glycol can be an aryl or alkylene glycol, preferably 20 selected from the group of glycerol, polyethylene glycol, panthanol, and sorbitol. If the glycol is an alkylene glycol it is preferably propylene glycol present at from about 10.0% to 20.0% by weight.

The water used in the gel is preferably purified and 25 pyrogen free water and is present in an amount sufficient to bring the total composition up to 100% by weight.

Various optional ingredients can also be included in the gel composition such as preservatives eg, methylhydroxybenzoate and propylhydroxybenzoate. In 30 addition, the wound gel composition can, if desired, contain small amounts (effective amounts) i.e. less than

5 5%, of pharmacologically active ingredients. For example,
an antibiotic or antimicrobial agent such as metronidazole,
silver sulphadiazine, neomycin or penicillin, and
antiseptic agent such as povidone iodine, and anti-
inflammatory agent such as hydrocortisone or triamcinolone
10 acetonide, or a skin protective agent such a zinc oxide can
be included.

We have found that the invention performs particularly well
if the gel has a viscosity of from 150 to 800 Pas as
determined by Viscolog model MRV8 viscometer and a helical
15 drive unit and PD spindal rotating at 2.5 rpm. Gels
packaged in this way have been found to have particularly
homogeneous viscosity due to the uniformity of the package
compared to that found in tubes or other less symmetrical
dispensing devices.

20 The barrier vessel containing a wound gel of the invention
may be made by the method described in EP 0418724 or EP
0017147 to Lechner GmbH. The vessel may be sealed by a
valve having a cup and discharge port, particularly of the
type CA38F/39F ex Rexam Dispenser, Portsmouth UK, although
25 valves having a gasket able to withstand steam
sterilisation would be suitable for producing a sterile
product.

We have also found that the rate at which gel is dispensed
may be altered by altering the applicator nozzle size.
30 Thus it is possible to change the applicator in order to
get fast release or slow release of the gel which may be
important for some wounds.

- 5 The invention is illustrated by the following non-limiting examples.

Example 1

	<u>Gel composition</u>	<u>% by weight</u>
	Pectin	0.1%
10	Sodium carboxymethyl cellulose	3.4%
	Propylene glycol	15.0%
	Purified water	81.5%

Pectin (0.2g) was added to purified water (163.0g) in a beaker and heated to 50 - 60°C with constant stirring until
15 the pectin dissolved. Propylene glycol (30.0g) was added and sodium carboxymethyl cellulose (6.8g) was gradually added with constant mixing. A hydrocolloid gel (200g) was produced.

Example 2

- 20 The gel from example 1 was used to fill the inner container of a barrier aerosol vessel and a valve having a cup and discharge port applied to seal the vessel. The filled container was terminally steam sterilised for 30 minutes at 121C. The vessel was then removed to a clean room and an
25 applicator fitted to the valve and the outer container gassed to pressurise the gel.

Example 3

The barrier aerosol vessel containing gel prepared as in

5 Example 2 was subjected to a microbial challenge. A mixed
microbial suspension (S.aureus, E.coli and C. albicans -
all typical wound bacteria) was prepared at a concentration
of 1×10^5 /ml and inoculated into the first 2cms of gel
contained in the nozzle of the vessel. The inoculated
10 canned gel was then left to stand at room temperature for
a period of 7 days and then sampled. This mimics clinical
use. After sampling the gel was re-inoculated with the
microbial suspension and sampled after a further 7 days.
Sampling was achieved by 10 fold dilution plating out
15 appropriate dilutions onto pre-dried TSA plates. The
plates were incubated at 35C for 24/48 hours prior to
counting.

The results of the assay demonstrated a 5 log reduction in
each of the three challenge organisms over days 0-7 and 7-
20 14. These results show that micro-organisms do not
proliferate in the gel contained in the barrier vessel.
This makes the combination of gel and barrier vessel
suitable for a multi-dose sterile product.

1 / 1

FIG. 1





FIG. 2



THIS PAGE BLANK (USPTO)